

Exhibit 91

Academy Press Tuesday, Jan. 16, 1996 B5

Condom makers stop using talc

By MARIE MC GULLOUGH
KNIGHT-RIDDER NEWSPAPERS

Candace Sue Kasper believes "safe sex" should be as safe for women as for men.

So early this year, the Dallas skin pathologist began urging — some would say badgering — condom makers and the federal Food and Drug Administration to stop the little-known practice of coating condoms with talc.

Talc, a powder made from the rock-like mineral magnesium silicate, is an excellent dry lubricant, but can scar soft tissues inside the body, where it does not dissolve. In women, body powders containing talc have been linked to infertility and ovarian cancer.

Kasper's campaign apparently worked.

"We've requested U.S. manufacturers to cease using (talc) and, in fact, all have agreed not to use it in manufacturing condoms," FDA spokesman Arthur Whitmore said last month.

Carter-Wallace, which makes Trojans and claims 60 percent of the American condom market, said in a statement that "to allay any possible concern," it has "discontinued the use of talc in its condom manufacturing process."

Kasper, 46, feels vindicated, but not victorious. She said an FDA official told her the agency had no rules against talc on condoms and no way to check compliance.

"When we have someone independent policing it, that's when I'll feel I have a victory," Kasper said.

Sales of condoms increased nearly 50 percent in the late 1980s, as AIDS awareness grew, according to the U.S. Centers for Disease Control and Prevention.

Kasper, an advocate of condom use, applauds the trend. And she is the first to say no research, including her own, has proved that talc specifically from condoms causes reproductive health problems in women.

Concern about talc as an ovarian carcinogen goes back 50 years in the medical literature. By the 1970s, evidence was mounting that talc particles might migrate into a woman's fallopian tubes where they could cause scarring and irritation of the ovaries. Scientists believed in some cases that the scarring led to infertility or cancer.

In 1992, Bernard Harlow, an epidemiologist and obstetrician-gynecol-

ogist at Harvard Medical School, published a study that found that women who used powder on their genitals, undergarments or diaphragms over a long period of time increased their risk of ovarian cancer threefold. (Today, feminine hygiene products use ingredients other than talc, though many baby and dusting powders still contain talc.)

Like Kasper, Harlow believes there is no reason to put talc on condoms — even if it has not been proven to be harmful — because cornstarch is a cheap, safe alternative.

"We'll probably never know for sure" that condom talc is unsafe, he said. "But why take the risk? Cornstarch does just fine and doesn't pose risks. I think it's prudent for manufacturers" to switch.

Indeed it is, said Ansell Inc. of Easton, Pa., which makes LifeStyle condoms and has about a quarter of the American condom market. Ansell switched from talc to cornstarch in January 1994.

"We knew surgical glove talc was a problem, so we figured there might be a problem with condoms," said Milt Hinech, Ansell's vice president of technical affairs. "Whether it's rational or

scientific, you just have to say, 'Let's not argue about it. Let's just do it.'"

In 1990, the FDA asked manufacturers to voluntarily stop putting talc on surgical gloves amid mounting scientific evidence that it caused adhesions in surgical patients. At the same time, the agency evaluated talc on condoms, but concluded the amount was insignificant and did not pose problems, said FDA spokeswoman Sharon Snider.

Kasper's concern about condoms arose after she and a colleague, Dallas plastic surgeon Preston Chandler, discovered talc in the hardened tissue surrounding breast implants that had been removed from women.

In a 1994 journal article, the two physicians speculated that the talc might have contributed to the hardening and that it came from surgical gloves. They also speculated that talc might play a role in the autoimmune symptoms that are the subject of numerous breast implant lawsuits.

Curious to see whether other products were dusted with talc, Kasper and Chandler bought condoms, pacifiers and baby-bottle nipples in 1994 at Dallas-area stores, then scrutinized

them under a microscope. The nipples and pacifiers appeared clean, but all eight brands of American-made and foreign-made condoms had varying amounts of talc, cornstarch and, in some cases, substances such as sand, silicone dioxide or club moss spores (an outmoded lubricant that also causes scarring in soft tissues).

Kasper and Chandler wrote to the condom manufacturers, several of which responded that they did not use talc in their production process. Carter-Wallace did not respond to them, Kasper said.

Only one manufacturer, Ansell, backed up its claim to be talc-free, Kasper said. She examined Ansell condoms made after January 1994 and found cornstarch, not talc.

She and Chandler also expressed their concern to the FDA, which thanked them for their information — but didn't say it would take any action.

"Largely, we've been ignored," said Kasper, who has a private pathology practice and is a staff physician at Baylor Medical Center. "Fortunately, my livelihood doesn't depend on this. I've done this on my own time and money."



AMERICAN HEALTH CONSULTANTS

CONTRACEPTIVE TECHNOLOGY

U P D A T E

A Monthly Newsletter for Health Professionals

INSIDE

■ How to pick which condoms to distribute ... 135

■ What's the problem with talc? 136

■ So, you want to spread the word about ECPs? Make sure you're not breaking the law 137

■ Case studies to help you treat older women 142

■ *CTU* updates: Get your copy of the CDC's new report on critical health concerns of women, children 144

■ Inserted in this issue: Results from *CTU's* 1995 exclusive Salary Survey

**NOVEMBER
1995**

VOL. 16 NO. 11
(pages 133-144)

American Health Consultants is
A Medical Economics Company

You can steer clients to condoms free from potentially harmful talc

Condom companies agree to produce without the dry lubricant

Due to reports earlier this year that talc on condoms potentially could lead to infertility and ovarian cancer, some clinicians began warning clients to use only brands known to be talc-free. *Contraceptive Technology Update* recently contacted the major condom companies to determine if plant manufacturing processes had changed to eliminate talc, and all responded that, indeed, there is no longer talc on their condoms. They and other experts give you advice on how to counsel patients about finding talc-free condoms (see box, p. 135).

Some companies told us they didn't use talc at the time a letter to the editor of the *Journal of the American Medical Association (JAMA)* indicated the silicate had been found on their brands.¹ The authors of the letter stand behind their work and have provided *CTU* with copies of all correspondences between them and the companies, as well as with the U.S. Food and Drug Administration (FDA) in Rockville, MD.

Through the use of polarization microscopy, Candace Sue Kasper, MD, PhD, director of Southwest Dermatopathology Consultants in Dallas, and her colleague P.J. Chandler, MD, MBA, owner of Ambulatory Plastic Surgery Center, also in Dallas, showed that "maximum density" of talc particles could be found on the following:

- TrojanENZ nonlubricated condoms (Carter Wallace, Cranberry, NJ);
- Kimono lubricated condoms (Sagami Rubber, Tokyo/Mayer Laboratories, Oakland, CA);

- RIA lubricated condoms (Summa Trading Co., Malaysia).

They found "few to numerous" particles of talc in:

- Sagami lubricated condoms (Sagami Rubber/Mayer Laboratories);
- Lifestyle lubricated condoms (Ansell Consumer Products, Eatontown, NJ).

And they found "rare to few" particles in:

- Ramses Ultrathin condoms (London International U.S. Holdings, Sarasota, FL);

- Gold Circle Coin condoms (Aladan Corp., Norcross, GA).

The polarization examination, conducted with crossed filters that

block out light from all particles except those that are capable of bending light — like talc — was conducted during 1994. In addition to sending the results to *JAMA* at the end of 1994, Kasper sent them to the condom manufacturers and the FDA, she says. The letter appeared in *JAMA* on March 15, 1995.

David Mayer, president of Mayer Labs, a company that imports and distributes condoms in the United States, expressed surprise that talc had been found on the condoms he sells. He wrote to Kasper in January 1995 asking her to remove his condoms from the list of talc-containing products. Mayer attached a correspondence from Minoru Takahashi, director of the condom production division of Sagami Rubber, which makes Mayer's condoms under contract. Takahashi writes, "Sagami assures you that we never use the talc on all condoms manufactured by us as well as Kimono and Maxx series condoms to Mayer Laboratories."

After the *JAMA* letter appeared with Sagami/Mayer condoms still listed, Mayer wrote Kasper once again in May, stating, "... we require that you write a subsequent letter to *JAMA* informing the editor of the inaccuracy of your data and retracting the names of our products ..."

A "puzzled" Kasper responded in a letter that "... because we have confidence in our identification techniques, we see no reason to retract our findings. On the contrary, we think that it is you who needs to respond to us with full disclosure of the identity of all substances used in the manufacture and packaging of your condoms."

A more public exchange — letters from Mayer and Kasper — is due soon in *JAMA* but had not appeared at press time.

It has been almost 10 years since talc was used in the manufacturing of his company's condoms, Mayer tells *CTU*. He suggests the crystalline particles seen by Kasper are some other substance and not talc, but Kasper is dubious.

"There are some other crystals they might be, but it's unlikely," she says of the needle-like microscopic structures.

FDA: No talc on American condoms

After contacting the FDA in late 1994, Kasper says she was referred to the agency's division of compliance, where she learned that the condom companies could not be considered out of compliance because there were no guidelines about talc on condoms. Even if there were guidelines, they

would apply only to new condoms, she was told.

Raju Kammula, PhD, DVM, the FDA's chief of toxicology, doesn't think the lack of guidelines is a problem since condom manufacturers are market-driven.

"The manufacturers comply voluntarily a lot of the time," he says. "They realize that if there's a problem and it's not corrected, their products won't be sold. I think this is one incentive they have. They want to promote a good product, and if there is any evidence that it is going to cause problems, their business will be affected."

Besides, guidelines aren't always necessary, since the site inspectors are trained to evaluate the manufacturing process, and talc is "something they can zero in on in a fairly straightforward manner," says Colin Pollard, chief of the FDA's OB/GYN devices branch. "Talc is an obvious part of the manufacturing process. It would be easy [for inspectors] to see." The inspectors also view "master files" of the manufacturing process kept by the companies and would notice talc immediately, he adds.

Site inspectors don't constantly keep tabs on plants in Japan and other foreign countries, but Pollard says the four U.S. manufacturers — London International (condoms are processed in Anderson, SC), Carter Wallace, Aladan, and Ansell — account for nearly 99% of the American market. All but Carter Wallace had indicated they stopped using talc prior to being contacted by the FDA, he says.

Ansell stopped using talc in January 1994 and began using a starch powder, according to a letter to Kasper from the company's vice president of technical affairs, J.W. Moushall. Moushall pointed out that talc levels prior to that were below 1% of all particles on the condom, based on results from Federal Bureau of Investigation (FBI) investigations of sex offenders who use condoms to avoid detection through semen analysis.

Carter Wallace last to switch

The FDA had subsequent discussions with Carter Wallace, "and as a result of that exchange, the company decided to take talc out of the [manufacturing] process," says Pollard.

One reason the company may have been reluctant to switch away from talc is because scientists there believed talc did not cause health problems. Based on presentations at a workshop co-sponsored by the FDA and the Columbia, MD-based International Society of Regulatory

Distribute the correct condoms at your facility

Now that you know the potential problems caused by talc on condoms, what should you do about it? Researchers who have been studying the possible health risks say you should simply choose the right condoms to distribute to your clients.

Candace Sue Kasper, MD, PhD, director of Southwest Dermatopathology Consultants in Dallas, is adamant that she doesn't want people to stop using condoms. Kasper simply wants condom manufacturers to use a substitute product that will work just as well as talc, such as corn starch. Family planning practitioners agree.

"It seems to me, if you have a choice, and one of the choices has unknown health risks, why use it?" asks Dolly Joern, ARNP, a clinician at Mount Vernon (WA) Women's Clinic.

Joern read reports linking condom use and talc exposure and began talking with patients about it. A "very relevant" topic for her since her mother and aunt both died of ovarian cancer, Joern has been suggesting to her clients that they use Lifestyles condoms.

Find companies that have announced their

manufacturing system is free of talc and offer that company's condoms to clients, Kasper suggests. Lifestyles are a good choice, since the company was the first to acknowledge the need to switch to a different lubricant, she says.

All U.S. condom manufacturers have reportedly switched to other lubricants, such as corn starch or other types of food starch, says Raju Kammula, PhD, DVM, the Food and Drug Administration's chief of toxicology. Condoms made after March 1995 should be free from talc, he says.

The only problem that might arise is using condoms manufactured before the companies agreed to change their process. David Mayer, president of Mayer Labs, an Oakland, CA-based company that imports and distributes Kimono and Maxx brand condoms in the United States, also made two suggestions for clinicians and their patients. Since condoms manufactured before companies switched to new lubricants might still contain talc, he recommends checking all condom expiration dates. Choose ones with the dates most in the future; the normal expiration time is five years, so dates after 2000 should be free of talc.

When buying condoms in stores, shop where there would be a large volume of condoms purchased, Mayer says. This will ensure that the supplies are as fresh as possible. ■

Toxicology and Pharmacology in January 1994 called, "Talc: Consumer Uses and Health Perspectives," Carter Wallace issued a statement pointing out that:

- "studies . . . indicate the trace amounts of talc on a condom purchased by the consumer should present no known health concern";
- "the workshop concluded that when taken together, the results of these studies are insufficient to demonstrate any real association" between talc and ovarian tumors;
- ". . . there is no basis to conclude that talc is capable of migrating to the ovaries in the first place . . ."

"That's what Carter Wallace wanted to come away with from [the workshop]. I didn't necessarily agree with that finding," says Bernard L. Harlow, PhD, a researcher in the OB/GYN epidemiology center of Brigham and Women's Hospital at Harvard Medical School in Boston. He presented a review of the scientific literature

related to perineal talc exposure and the risk of ovarian cancer, which has recently been published.²

While Harlow says talc exposure may not be as big a risk factor for ovarian cancer as it once was, since women tend to use less talc applied directly to the perineum, new exposures — from condoms, for example — could change that scenario, he says.

The range of relative risk estimates in various studies of the association between ovarian cancer and perineal talc exposure is between 1.0 and 1.8,¹ which is plausible, says Harlow (1.0 would indicate no association; 2.0 would indicate a clear association).

"Many other risk factors contribute to the development of ovarian cancer much more so than perineal exposure to talc," he adds.

When it was found that tubal ligation was protective against ovarian cancer, researchers speculated it might be due to the inability of a carcinogen to travel up the tubes to the ovaries.³ But the

Why is talc considered a problem substance?

Talc is a natural mineral, hydrous magnesium silicate in the asbestos family. Although it is a known sclerosing agent^{1,2} and has been implicated as a lung carcinogen^{3,4} and an ovarian carcinogen,^{5,7} it is a prevalent substance in everyday life and can be found in everything from aspirin to cosmetics, says Candace Sue Kasper, MD, PhD, director of Southwest Dermatopathology Consultants in Dallas.

The "good, cheap lubricant" is safe as long as it doesn't come into contact with open tissue lesions, Kasper says. It is often used as a dry lubricant by the condom industry to facilitate the removal of dipped latex from the molds for foiling and filming packaging machines, and to keep the rolled condoms from sticking to each other before they can be packaged.

"As pathologists, we're very familiar with talc. It's not anything that's a mystery. We know what it looks like, we know what it does," says Kasper. "So talc is OK. But once it gets into the soft tissue, then it can cause a problem."

Talc doesn't harm people when it is ingested (unless they have ulcers or other open lesions in the digestive system), because peristalsis pushes it through the system and it has an exit. Therein lies the problem with the lungs and the ovaries.

"The difference with the ovary is that anything that gets up through the fallopian tube and passes to the ovary does not have an exit. There's nowhere for it to go," Kasper explains.

Talc can travel up the reproductive tract and sit on the surface of the ovary, she believes. When the ovary opens to release an egg, talc can get trapped inside. When the body mounts a defense and the talc is consumed by macrophages, cytokines are produced. It is

these cytokines that can produce the fibrosis and scarring — and therefore infertility — talc has been associated with, Kasper says.

Kasper and her colleague P.J. Chandler, MD, MBA, of Ambulatory Plastic Surgery Center, also in Dallas, became interested in talc in 1989 when they were trying to determine why women's silicone breast implants sometimes made their breasts hard. At the time, Kasper worked at the University of Texas Southwestern Medical School in Dallas. A clinicopathologic study of the fibrotic capsules surrounding the implants turned up talc on 70% of the capsules,⁸ probably from latex surgical gloves manufactured prior to 1991.¹ Curious about other latex products that might contain talc, they systematically purchased condoms, dental dams, baby bottle nipples, and pacifiers from local stores and began analyzing them under the microscope. While the baby products contained no talc, the condoms and the dental dams often did.¹

References

1. Kasper CS, Chandler PJ. Possible morbidity in women from talc on condoms. *JAMA* 1995; 273:846-847.
2. Saxen L, Kassinen A, Saxen E. Peritoneal foreign-body reaction caused by condom emulsion. *Lancet* 1963; 1:1295-1296.
3. Thomas TL, Stewart PA. Mortality from lung cancer and respiratory disease among pottery workers exposed to silica and talc. *Am J Epidemiol* 1987; 14:35-43.
4. Kleinfeld M, Messite J, Zaki MH. Mortality experience among talc workers: A follow-up study. *J Occup Med* 1974; 16:345-349.
5. Longo DL, Young RC. Cosmetic talc and ovarian cancer. *Lancet* 1979; 2:1011-1012.
6. Cramer DW, Welch WR, Scully RE, Wojciechowski CA. Ovarian cancer and talc. *Cancer* 1982; 50:372-376.
7. Harlow BL, Cramer DW, Bell DA, et al. Perineal exposure to talc and ovarian cancer risk. *Obstet Gynecol* 1992; 80:19-26.
8. Kasper C, Chandler P. Talc deposition in skin and tissues surrounding silicone gel-containing prosthetic devices. *Arch Dermatol* 1994; 130:48-53. ■

121,700 study subjects were asked about talc use on sanitary napkins or directly on the perineum. When analyzed, the relationship between talc exposure and ovarian cancer was not statistically significant, principal investigator Susan Hankinson, ScD, a researcher at Channing Laboratory in Boston, told CTU last year when her study was published.

(See CTU, May 1994, pp. 67-70.)

It would be difficult to compare perineal exposure and exposure from a condom, and even more difficult to measure talc exposure from condoms, Harlow says. Retrospectively, study subjects have trouble remembering how often they used condoms, much less if it was a brand that

had talc on it; a prospective study could not ethically expose a group of people to a substance that was potentially harmful.

Scientists at the Centers for Disease Control and Prevention (CDC) in Atlanta soon may be able to shed some light on the issue, however. During a recent preliminary review of an existing data set, no evidence for an association between those who had ever used a condoms and the risk for ovarian cancer could be found, says Bert Peterson, MD, chief of the women's health and fertility branch. Publication is pending, so details aren't available, but Peterson says the data set is large enough to generate meaningful evidence.

More studies needed

After conducting his extensive review, which he calls "the definitive paper on what we know about talc and where we need to go," Harlow concluded that more studies should be conducted to confirm or deny reports that talc can migrate and embed in human ovarian tissue. He was surprised to discover only two studies that investigated the issue, he says.

Kasper agrees more studies are needed. She would like to continue analyzing condoms for talc and other contaminants, but has left her position at the University of Texas Southwestern Medical School in Dallas and currently doesn't have easy access to the equipment she would need, she says. She did recheck a Lifestyles condom this summer, and it was free of talc.

She also recently rechecked a Ramses condom (which didn't have much talc to begin with) which still had high concentrations of lycopodium, a dry lubricant that comes from the spores of club moss, part of the fern family. In a statement to CTU, London International's Vice President of Marketing John Blutenthal writes: "London International U.S. Holdings Inc., makers of Ramses, Sheik, Touch, Avanti, and Fourex condoms, does not use talc or lycopodium in the manufacturing of its condoms."

References

1. Kasper CS, Chandler PJ. Possible morbidity in women from talc on condoms. *JAMA* 1995; 273: 846-847.
2. Harlow BL, Hartge PA. A review of perineal talc exposure and risk of ovarian cancer. *Reg Toxicol Pharmacol* 1995; 21:254-260.
3. Hankinson SE, Hunter DJ, Colditz GA, et al. Tubal ligation, hysterectomy, and risk of ovarian cancer. *JAMA* 1993; 270:2813-2818. ■

What are legalities of promoting ECPs?

Bottom line: It's 'defensible but risky'

Women should have wider access to emergency contraceptive pills (ECPs), an international group of experts recommended recently. But since the regimen involves using an approved drug for an unlabeled purpose, some providers worry they may be breaking the law by advertising or marketing the availability of the oral contraceptive (OC) pills for that purpose. Worse yet, some clinicians may be moving forward with highly visible marketing campaigns without knowing they are making themselves vulnerable to a battle in the courts.

Sometimes called postcoital or morning-after contraception, emergency methods include regular combined ethinyl estradiol/levonorgestrel pills given at a higher dose (the Yuzpe regimen) than normally prescribed and the copper IUD.

Because the legal boundaries about advertising emergency contraception have never been clear (see *Contraceptive Technology Update*, March 1993, pp. 33-45), CTU spoke with several lawyers to find out why. We learned that the basis for the law prohibiting the advertisement of unlabeled drugs is the 1938 Food, Drug, and Cosmetic Act and that the Food and Drug Administration (FDA) in Rockville, MD, does have authority to bring action against those who break the law. The FDA sets priorities regarding what issues will get the agency's attention, however.

In April, 24 experts representing the fields of research, policy, communications, women's advocacy, and medicine met in Bellagio, Italy, to discuss emergency contraception and made several recommendations. (See related article, p. 139.) They also ascertained the three reasons emergency contraception is not more widely available. They are:

- Women and providers are uninformed about the methods.
- Few products are marketed for emergency contraceptive use.
- Service providers are too often reluctant to provide the methods.

The meeting was held because many health care providers don't have an understanding of this important method of pregnancy prevention, says Elizabeth Robinson, associate director of

1. Dietrich E, Cohan C. *Women and Heart Disease*. New York, NY: Ballantine Books; 1992.
2. Kaplan N, Lieberman E, Neal W. *Clinical Hypertension*. Baltimore, MD: Williams & Wilkins Inc.; 1994.
3. Wild R, Taylor E, Knehans A. The gynecologist and the prevention of cardiovascular disease. *Am J Obstet Gynecol* 1995; 172:1-13.
4. 1988 Joint National Committee. The 1988 report of the joint national committee on detection, evaluation, and treatment of high blood pressure. *Arch Internal Med* 1988; 148:1023-1038.
5. Manson JE, Tosteson H, Ridker PM, et al. The primary prevention of myocardial infarction. *N Engl J Med* 1992; 326:1406-1416. ■

Correction

In the September 1995 issue of *Contraceptive Technology Update*, the continuing education objectives inadvertently were omitted. Following are the continuing education nursing objectives for that issue:

1. Describe the reasons triphasic pills are prescribed in the clinic setting.
2. Cite the issues surrounding the over-the-counter oral contraceptive movement.
3. Describe the social and cultural effects of the Pill.
4. Counsel pill users about their risk for breast cancer.


CTU regrets the error. ■

Continuing education objectives

After reading this issue of *Contraceptive Technology Update*, the reader will be able to:

1. Describe the various Norplant implant removal techniques.
2. Find necessary resources to prepare for Norplant removals.
3. Cite several ways to better perform Norplant removals.
4. Describe how to counsel contraceptive patients about their risks for cardiovascular disease. ■

Contraceptive Technology Update is endorsed by the National Association of Nurse Practitioners in Reproductive Health and the Association of Reproductive Health Professionals as a vital information source for health care professionals.



A • R • H • P
Association of Reproductive Health
Professionals

EDITORIAL ADVISORY BOARD

Chairman:

Robert A. Hatcher, MD, MPH
Senior Author, *Contraceptive Technology*
Professor of OB/GYN and Director, Emory University/Grady Memorial
Hospital Family Planning Program
Atlanta

David F. Archer, MD
Professor of OB/GYN
The Jones Institute for
Reproductive Medicine
Norfolk, VA

Deanis J. Barbour, JD
President, Association of Reproductive
Health Professionals
Washington, DC

Wills Brown, MD, MPH
Director, Personal Health Services
Howard County Health Dept.
Columbia, MD

Joyce Carlone, RNC, MN
Director of GYN/Ob
Nurse Practitioner Program
Emory University Regional Training Center for
Family Planning
Atlanta

Andrew M. Kunitz, MD
Professor of OB/GYN
Director, Division of Ambulatory Care
Univ. of Florida Health Sciences Center
Jacksonville, FL

Anita Nelson, MD
Medical Director,
Women's Health Care Clinic Harbor/UCLA
Medical Center
Torrance, CA

Michael Rosenberg, MD, MPH
Clinical Associate Professor of OB/GYN and
Epidemiology
University of North Carolina
President, Health Decisions Inc.
Chapel Hill, NC

Allan Rosenbloom, MD
Dean, School of Public Health, Columbia
University
New York City

Sharon E. Schwab, RN, FNP, CNM, MSN
Family Planning Clinician and Consultant
Seattle

William Spillney, MD
Professor and Chairman,
Department of OB/GYN
University of South Florida
College of Medicine
Tampa, FL

James Trussell, PhD
Professor of Economics
and Public Affairs
Director, Office of
Population Research
Associate Dean, Woodrow Wilson School of
Public and
International Affairs
Princeton University
Princeton, NJ

Susan Wysocki, RNC, BSN, NP
Executive Director
National Association of Nurse Practitioners in
Reproductive Health
Chair, National Nurse
Practitioner Coalition
Washington, DC

Contraceptive Technology Update (ISSN 0274-728X), including *Women's Health Update* and *STD Quarterly*, is published monthly by American Health Consultants, 3525 Piedmont Road, Building Six, Suite 400, Atlanta, GA 30304. Telephone: (404) 252-7436. Second-class postage paid at Atlanta, GA 30304. POSTMASTER: Send address changes to *Contraceptive Technology Update*, P.O. Box 740059, Atlanta, GA 30374.

This continuing education offering is sponsored by American Health Consultants, which is accredited as a provider of continuing education in nursing by the American Nurses Credentialing Center's Commission on Accreditation. American Health Consultants is accredited by the Accreditation Council for Continuing Medical Education to sponsor CME for physicians. American Health Consultants designates this continuing medical education activity for 18 credit hours in Category 1 of the Physicians' Recognition Award of the American Medical Association. This CME activity was planned and produced in accordance with the ACCME Essentials. Physician members of American Health Consultants' 1995 Continuing Medical Education Council: Anthony Vuturs, MD, MPH; Stephen A. Brunton, MD; Dan Longo, MD; Lora Denton, MD; and Gideon Bosker, MD.

Opinions expressed are not necessarily those of this publication. Mention of products or services does not constitute endorsement. Clinical, legal, tax, and other comments are offered for general guidance only; professional counsel should be sought for specific situations.

Editor: Virginia Mason,
(404) 667-0073

President: Robert Williford,
(404) 262-7758, ext. 157.

Publisher: Connie Austin,
(404) 262-7759, ext. 152.

Executive Editor: Lisa Gandy Wargo,
(404) 262-7758, ext. 137.

Managing Editor: Joy Daugherty,
(404) 262-7769, ext. 140.

Production Editor: Nancy Baltmarsh,
(404) 262-7759, ext. 172.

Marketing Coordinator: Jennifer Lewis,
(404) 262-7759, ext. 207.

Copyright © 1995 by American Health Consultants. *Contraceptive Technology Update*, *Women's Health Update*, and *STD Quarterly* are trademarks of American Health Consultants. The trademarks *Contraceptive Technology Update*, *Women's Health Update*, and *STD Quarterly* are used herein under license. All rights reserved.

Hours of operation:
8:30 a.m. - 4:30 p.m.

Editorial Questions

For questions or comments, call Joy
Daugherty at (404) 262-7759, ext. 140.

Subscriber Information

Customer Service: (800) 888-2421

Subscription rates: U.S.A., one year (12 issues), \$289. With 18 nursing contact hours or Category 1 CME credits. \$339: Outside U.S., add \$20 per year, total prepaid in U.S. funds. One to nine additional copies, \$99 per year; 10 or more additional copies, \$75 per year. Call for more details. Back issues, when available, are \$25 each (GST registration number R128870672). Photocopying: No part of this newsletter may be reproduced in any form or incorporated into any information retrieval system without the written permission of the copyright owner. For reprint permission, please contact Lawrence Cohen at American Health Consultants. Address: P.O. Box 740056, Atlanta, GA 30374. Telephone: (800) 446-1139, ext. 214. Fax: (800) 753-9151.

Chudkowski, Michael

From: Christensen, Catharine
To: Chudkowski, Michael
Subject: Talc article
Date: Monday, October 23, 1995 9:28AM

October 23, 1995

WHY IS TALC CONSIDERED A PROBLEM SUBSTANCE?

Contraceptive Technology Update via First: Talc is a natural mineral, hydrous magnesium silicate in the asbestos family. Although it is a known sclerosing agent^{1,2} and has been implicated as a lung carcinogen^{3,4} and an ovarian carcinogen,⁵⁻⁷ it is a prevalent substance in everyday life and can be found in everything from aspirin to cosmetics, says Candace Sue Kasper, MD, PhD, director of Southwest Dermatopathology Consultants in Dallas.

The "good, cheap lubricant" is safe as long as it doesn't come into contact with open tissue lesions, Kasper says. It is often used as a dry lubricant by the condom industry to facilitate the removal of dipped latex from the molds for rolling and filming packaging machines, and to keep the rolled condoms from sticking to each other before they can be packaged.

"As pathologists, we're very familiar with talc. It's not anything that's a mystery. We know what it looks like, we know what it does," says Kasper.

"So talc is OK. But once it gets into the soft tissue, then it can cause a problem."

Talc doesn't harm people when it is ingested (unless they have ulcers or other open lesions in the digestive system), because peristalsis pushes it through the system and it has an exit. Therein lies the problem with the lungs and the ovaries.

"The difference with the ovary is that anything that gets up through the fallopian tube and passes to the ovary does not have an exit. There's nowhere for it to go," Kasper explains.

Talc can travel up the reproductive tract and sit on the surface of the ovary, she believes. When the ovary opens to release an egg, talc can get trapped inside. When the body mounts a defense and the talc is consumed by macrophages, cytokines are produced. It is these cytokines that can produce the fibrosis and scarring - and therefore infertility - talc has been associated with, Kasper says.

Kasper and her colleague P.J. Chandler, MD, MBA, of Ambulatory Plastic Surgery Center, also in Dallas, became interested in talc in 1989 when they were trying to determine why women's silicone breast implants sometimes made their breasts hard. At the time, Kasper worked at the University of Texas Southwestern Medical School in Dallas. A clinicopathologic study of the fibrotic capsules surrounding the implants turned up talc on 70% of the capsules,⁸ probably from latex surgical gloves manufactured prior to 1991.¹ Curious about other latex products that might contain talc, they systematically purchased condoms, dental dams, baby bottle nipples, and pacifiers from local stores and began analyzing them under the microscopes. While the baby products contained no talc, the condoms and the dental dams often did.¹


References

1. Kasper CS, Chandler PJ. Possible morbidity in women from talc on condoms. *JAMA* 1995; 273:846-847.
2. Saxen L, Kassinen A, Saxen E. Peritoneal foreign-body reaction caused by condom emulsion. *Lancet* 1963; 1:1295-1298.
3. Thomas TL, Stewart PA. Mortality from lung cancer and respiratory disease among pottery workers exposed to silica and talc. *Am J Epidemiol* 1987; 14:35-43.
4. Kleinfeld M, Messite J, Zeki MH. Mortality experience among talc workers: A follow-up study. *J Occup Med* 1974; 16:345-349.
5. Longo DL, Young RC. Cosmetic talc and ovarian cancer. *Lancet* 1979; 2:1011-1012.
6. Cramer DW, Welch WR, Scully RE, Wojciechowski CA. Ovarian cancer and talc. *Cancer* 1982; 50:372-376.
7. Harlow BL, Cramer DW, Bell DA, et al. Perineal exposure to talc and ovarian cancer risk. *Obstet Gynecol* 1992; 80:18-26.
8. Kasper C, Chandler P. Talc deposition in skin and tissues surrounding silicone gel-containing prosthetic devices. *Arch Dermatol* 1994; 130:48-53.



October 23, 1995

DISTRIBUTE THE CORRECT CONDOMS AT YOUR FACILITY



Contraceptive Technology Update via First! : Now that you know the potential problems caused by talc on condoms, what should you do about it? Researchers who have been studying the possible health risks say you should simply choose the right condoms to distribute to your clients.

Candace Sue Kasper, MD, PhD, director of Southwest Dermatopathology Consultants in Dallas, is adamant that she doesn't want people to stop using condoms. Kasper simply wants condom manufacturers to use a substitute product that will work just as well as talc, such as corn starch. Family planning practitioners agree.

"It seems to me, if you have a choice, and one of the choices has unknown health risks, why use it?" asks Dolly Joern, ARNP, a clinician at Mount Vernon (WA) Women's Clinic.

Joern read reports linking condom use and talc exposure and began talking with patients about it. A "very relevant" topic for her since her mother and aunt both died of ovarian cancer, Joern has been suggesting to her clients that they use Lifestyles condoms.

Find companies that have announced their manufacturing system is free of talc and offer that company's condoms to clients, Kasper suggests. Lifestyles are a good choice, since the company was the first to acknowledge the need to switch to a different lubricant, she says.

All U.S. condom manufacturers have reportedly switched to other lubricants, such as corn starch or other types of food starch, says Raju Kammula, PhD, DVM, the Food and Drug Administration's chief of toxicology. Condoms made after March 1995 should be free from talc, he says.

The only problem that might arise is using condoms manufactured before the companies agreed to change their process. David Mayer, president of Mayer Labs, an Oakland, CA-based company that imports and distributes Kimono and Maxx brand condoms in the United States, also made two suggestions for clinicians and their patients. Since condoms manufactured before companies switched to new lubricants might still contain talc, he recommends checking all condom expiration dates. Choose ones with the dates most in the future; the normal expiration time is five years, so dates after 2000 should be free of talc.

When buying condoms in stores, shop where there would be a large volume of condoms purchased, Mayer says. This will ensure that the supplies are as fresh as possible.

[10-21-95 at 11:46 EDT, Copyright 1995, American Health Consultants, File: b1019501.71w]

Picked by Profile: WHWomen's Healthcare

Source:
First! of INDIVIDUAL, Inc.

Contraceptive Technology Update via

Entire contents (C) 1995 by INDIVIDUAL, Inc., 84 Sherman Street, Cambridge, MA 02140.

First!

October 23, 1995

**YOU CAN STEER CLIENTS TO CONDOMS FREE FROM POTENTIALLY
HARMFUL TALC**

Contraceptive Technology Update via First! : Condom companies agree to produce without the dry lubricant

Due to reports earlier this year that **talc** on condoms potentially could lead to infertility and ovarian cancer, some clinicians began warning clients to use only brands known to be **talc**-free. Contraceptive Technology Update recently contacted the major condom companies to determine if plant manufacturing processes had changed to eliminate **talc**, and all responded that, indeed, there is no longer **talc** on their condoms. They and other experts give you advice on how to counsel patients about finding **talc**-free condoms (see box, p. 135).

Some companies told us they didn't use **talc** at the time a letter to the editor of the Journal of the American Medical Association (JAMA) indicated the silicate had been found on their brands.¹ The authors of the letter stand behind their work and have provided CTU with copies of all correspondences between them and the companies, as well as with the U.S. Food and Drug Administration (FDA) in Rockville, MD.

Through the use of polarization microscopy, Candace Sue Kasper, MD, PhD, director of Southwest Dermatopathology Consultants in Dallas, and her colleague P.J. Chandler, MD, MBA, owner of Ambulatory Plastic Surgery Center, also in Dallas, showed that "maximum density" of **talc** particles could be found on the following:

- TrojanENZ nonlubricated condoms (Carter Wallace, Cranberry, NJ);
- Kimono lubricated condoms (Sagami Rubber, Tokyo/Mayer Laboratories, Oakland, CA);
- RIA lubricated condoms (Summa Trading Co., Malaysia).

They found "few to numerous" particles of **talc** in:

- Sagami lubricated condoms (Sagami Rubber/Mayer Laboratories);
- Lifestyle lubricated condoms (Ansell Consumer Products, Eatontown, NJ).

And they found "rare to few" particles in:

- Ramses Ultrathin condoms (London International U.S. Holdings, Sarasota, FL);
- Gold Circle Coin condoms (Aladan Corp., Norcross, GA).

The polarization examination, conducted with crossed filters that block out light from all particles except those that are capable of bending light - like **talc** - was conducted during 1994. In addition to sending the results to JAMA at the end of 1994, Kasper sent them to the condom manufacturers and the FDA, she says. The letter appeared in JAMA on March 15, 1995.

David Mayer, president of Mayer Labs, a company that imports and distributes condoms in the United States, expressed surprise that talc had been found on the condoms he sells. He wrote to Kasper in January 1995 asking her to remove his condoms from the list of talc-containing

products. Mayer attached a correspondence from Minoru Takahashi, director of the condom production division of Sagami Rubber, which makes Mayer's condoms under contract. Takahashi writes, "Sagami assures you that we never use the talc on all condoms manufactured by us as well as Kimono and Maxx series condoms to Mayer Laboratories."

After the JAMA letter appeared with Sagami/ Mayer condoms still listed, Mayer wrote Kasper once again in May, stating, ". . . we require that you write a subsequent letter to JAMA informing the editor of the inaccuracy of your data and retracting the names of our products"

A "puzzled" Kasper responded in a letter that ". . . because we have confidence in our identification techniques, we see no reason to retract our findings. On the contrary, we think that it is you who needs to respond to us with full disclosure of the identity of all substances used in the manufacture and packaging of your condoms."

A more public exchange - letters from Mayer and Kasper - is due soon in JAMA but had not appeared at press time.

It has been almost 10 years since talc was used in the manufacturing of his company's condoms, Mayer tells CTU. He suggests the crystalline particles seen by Kasper are some other substance and not talc, but Kasper is dubious.

"There are some other crystals they might be, but it's unlikely," she says of the needle-like microscopic structures.

FDA: No talc on American condoms

After contacting the FDA in late 1994, Kasper says she was referred to the agency's division of compliance, where she learned that the condom companies could not be considered out of compliance because there were no guidelines about talc on condoms. Even if there were guidelines, they would apply only to new condoms, she was told.

Raju Kammula, PhD, DVM, the FDA's chief of toxicology, doesn't think the lack of guidelines is a problem since condom manufacturers are market-driven.

"The manufacturers comply voluntarily a lot of the time," he says. "They realize that if there's a problem and it's not corrected, their products won't be sold. I think this is one incentive they have. They want to promote a good product, and if there is any evidence that it is going to cause problems, their business will be affected."

Besides, guidelines aren't always necessary, since the site inspectors are trained to evaluate the manufacturing process, and talc is "something they can zero in on in a fairly straightforward manner," says Colin Pollard, chief of the FDA's OB/GYN devices branch. "Talc is an obvious part of the manufacturing process. It would be easy [for inspectors] to see." The inspectors also view "master files" of the manufacturing process kept by the companies and would notice talc immediately, he adds.

Site inspectors don't constantly keep tabs on plants in Japan and other foreign countries, but Pollard says the four U.S. manufacturers - London International (condoms are processed in Anderson, SC), Carter Wallace, Aladan, and Ansell - account for nearly 99% of the American market. All but Carter Wallace had indicated they stopped using talc prior to being contacted by the FDA, he says.

Ansell stopped using talc in January 1994 and began using a starch powder, according to a letter to Kasper from the company's vice president of technical affairs, J.W. Moushall. Moushall pointed out that talc levels prior to that were below 1% of all particles on the condom, based on results from Federal Bureau of Investigation (FBI) investigations of sex offenders who use condoms to avoid detection through semen analysis.

Carter Wallace last to switch

The FDA had subsequent discussions with Carter Wallace, "and as a result of that exchange, the

company decided to take talc out of the [manufacturing] process," says Pollard.

One reason the company may have been reluctant to switch away from talc is because scientists there believed talc did not cause health problems. Based on presentations at a workshop co-sponsored by the FDA and the Columbia, MD- based International Society of Regulatory Toxicology and Pharmacology in January 1994 called, "Talc: Consumer Uses and Health Perspectives," Carter Wallace issued a statement pointing out that:

- * "studies . . . indicate the trace amounts of talc on a condom purchased by the consumer should present no known health concern";
- * "the workshop concluded that when taken together, the results of these studies are insufficient to demonstrate any real association" between talc and ovarian tumors;
- * ". . . there is no basis to conclude that talc is capable of migrating to the ovaries in the first place"

"That's what Carter Wallace wanted to come away with from [the workshop]. I didn't necessarily agree with that finding," says Bernard L. Harlow, PhD, a researcher in the OB/GYN epidemiology center of Brigham and Women's Hospital at Harvard Medical School in Boston. He presented a review of the scientific literature related to perineal talc exposure and the risk of ovarian cancer, which has recently been published.²

While Harlow says talc exposure may not be as big a risk factor for ovarian cancer as it once was, since women tend to use less talc applied directly to the perineum, new exposures - from condoms, for example - could change that scenario, he says.

The range of relative risk estimates in various studies of the association between ovarian cancer and perineal talc exposure is between 1.0 and 1.8,¹ which is plausible, says Harlow (1.0 would indicate no association; 2.0 would indicate a clear association).

"Many other risk factors contribute to the development of ovarian cancer much more so than perineal exposure to talc," he adds.

When it was found that tubal ligation was protective against ovarian cancer, researchers speculated it might be due to the inability of a carcinogen to travel up the tubes to the ovaries.³ But the 121,700 study subjects were asked about talc use on sanitary napkins or directly on the perineum. When analyzed, the relationship between talc exposure and ovarian cancer was not statistically significant, principal investigator Susan Hankinson, ScD, a researcher at Channing Laboratory in Boston, told CTU last year when her study was published. (See CTU, May 1994, pp. 67-70.)

It would be difficult to compare perineal exposure and exposure from a condom, and even more difficult to measure talc exposure from condoms, Harlow says. Retrospectively, study subjects have trouble remembering how often they used condoms, much less if it was a brand that had talc on it; a prospective study could not ethically expose a group of people to a substance that was potentially harmful.

Scientists at the Centers for Disease Control and Prevention (CDC) in Atlanta soon may be able to shed some light on the issue, however. During a recent preliminary review of an existing data set, no evidence for an association between those who had ever used a condoms and the risk for ovarian cancer could be found, says Bert Peterson, MD, chief of the women's health and fertility branch. Publication is pending, so details aren't available, but Peterson says the data set is large enough to generate meaningful evidence.

More studies needed

After conducting his extensive review, which he calls "the definitive paper on what we know about talc and where we need to go," Harlow concluded that more studies should be conducted to confirm or deny reports that talc can migrate and embed in human ovarian tissue. He was

surprised to discover only two studies that investigated the issue, he says.

Kasper agrees more studies are needed. She would like to continue analyzing condoms for talc and other contaminants, but has left her position at the University of Texas Southwestern Medical School in Dallas and currently doesn't have easy access to the equipment she would need, she says. She did recheck a Lifestyles condom this summer, and it was free of talc.

She also recently rechecked a Ramses condom (which didn't have much talc to begin with) which still had high concentrations of lycopodium, a dry lubricant that comes from the spores of club moss, part of the fern family. In a statement to CTU, London International's Vice President of Marketing John Blutenthal writes: "London International U.S. Holdings Inc., makers of Ramses, Sheik, Touch, Avanti, and Fourx condoms, does not use talc or lycopodium in the manufacturing of its condoms."

References

1. Kasper CS, Chandler PJ. Possible morbidity in women from talc on condoms. JAMA 1995; 273: 846-847.
2. Harlow BL, Hartge PA. A review of perineal talc exposure and risk of ovarian cancer. Reg Toxicol Pharmacol 1995; 21:254-260.
3. Hankinson SE, Hunter DJ, Colditz GA, et al. Tubal ligation, hysterectomy, and risk of ovarian cancer. JAMA 1993; 270:2813-2818.

[10-21-95 at 11:45 EDT, Copyright 1995, American Health Consultants, File: b1019501.6(w)]

Picked by Profile: WHWomen's Healthcare

Source:

Contraceptive Technology Update via

First of INDIVIDUAL, Inc.

Entire contents (C) 1995 by INDIVIDUAL, Inc., 84 Sherman Street, Cambridge, MA 02140.



October 23, 1995

WHY IS TALC CONSIDERED A PROBLEM SUBSTANCE?



Contraceptive Technology Update via First! : Talc is a natural mineral, hydrous magnesium silicate in the asbestos family. Although it is a known sclerosing agent^{1,2} and has been implicated as a lung carcinogen^{3,4} and an ovarian carcinogen,⁵⁻⁷ it is a prevalent substance in everyday life and can be found in everything from aspirin to cosmetics, says Candace Sue Kasper, MD, PhD, director of Southwest Dermatopathology Consultants in Dallas.

The "good, cheap lubricant" is safe as long as it doesn't come into contact with open tissue lesions, Kasper says. It is often used as a dry lubricant by the condom industry to facilitate the removal of dipped latex from the molds for foiling and filming packaging machines, and to keep the rolled condoms from sticking to each other before they can be packaged.

"As pathologists, we're very familiar with talc. It's not anything that's a mystery. We know what it looks like, we know what it does," says Kasper. "So talc is OK. But once it gets into the soft tissue, then it can cause a problem."

Talc doesn't harm people when it is ingested (unless they have ulcers or other open lesions in the digestive system), because peristalsis pushes it through the system and it has an exit. Therein lies the problem with the lungs and the ovaries.

"The difference with the ovary is that anything that gets up through the fallopian tube and passes to the ovary does not have an exit. There's nowhere for it to go," Kasper explains.

Talc can travel up the reproductive tract and sit on the surface of the ovary, she believes. When the ovary opens to release an egg, talc can get trapped inside. When the body mounts a defense and the talc is consumed by macrophages, cytokines are produced. It is these cytokines that can produce the fibrosis and scarring - and therefore infertility - talc has been associated with, Kasper says.

Kasper and her colleague P.J. Chandler, MD, MBA, of Ambulatory Plastic Surgery Center, also in Dallas, became interested in talc in 1989 when they were trying to determine why women's silicone breast implants sometimes made their breasts hard. At the time, Kasper worked at the University of Texas Southwestern Medical School in Dallas. A clinicopathologic study of the fibrotic capsules surrounding the implants turned up talc on 70% of the capsules,⁸ probably from latex surgical gloves manufactured prior to 1991.¹ Curious about other latex products that might contain talc, they systematically purchased condoms, dental dams, baby bottle nipples, and pacifiers from local stores and began analyzing them under the microscope. While the baby products contained no talc, the condoms and the dental dams often did.¹

References

1. Kasper CS, Chandler PJ. Possible morbidity in women from talc on condoms. JAMA 1995; 273:846-847.
2. Saxen L, Kassinen A, Saxen E. Peritoneal foreign-body reaction caused by condom emulsion.

Lancet 1963; 1:1295-1296.

3. Thomas TL, Stewart PA. Mortality from lung cancer and respiratory disease among pottery workers exposed to silica and talc. Am J Epidemiol 1987; 14:35-43.

4. Kleinfeld M, Messite J, Zakl MH. Mortality experience among talc workers: A follow-up study. J Occup Med 1974; 16:345-349.

5. Longo DL, Young RC. Cosmetic talc and ovarian cancer. Lancet 1979; 2:1011-1012.

6. Cramer DW, Welch WR, Scully RE, Wojciechowski CA. Ovarian cancer and talc. Cancer 1982; 50:372-376.

7. Harlow BL, Cramer DW, Bell DA, et al. Perineal exposure to talc and ovarian cancer risk. Obstet Gynecol 1992; 80:19-26.

8. Kasper C, Chandler P. Talc deposition in skin and tissues surrounding silicone gel-containing prosthetic devices. Arch Dermatol 1994; 130:48-53.

[10-21-95 at 11:47 EDT, Copyright 1995, American Health Consultants, File: b1019501.8lw]

Picked by Profile: WHWomen's Healthcare

Source:
First of INDIVIDUAL, Inc.

Contraceptive Technology Update via

Entire contents (C) 1995 by INDIVIDUAL, Inc., 84 Sherman Street, Cambridge, MA 02140.